

## REMARKS ON FIXED POINT THEOREMS ON STAR-SHAPED SETS

BY SEHIE PARK AND JAIHAN YOON

A large number of fixed point theorems without convexity conditions are available in analysis. Some of them are related to the concept of star-shapedness.

In this paper, we obtain a fixed point theorem which extends known results of Meir-Keeler [13], Boyd-Wong [5], Browder [6], and the Banach contraction principle. Using this, we extend and unify fixed point theorems of Assad [3], Assad-Kirk [1], Kuhfittig [11], and Dotson [8] on convex or star-shaped sets. Finally, related results of Rhoades [14] and Assad [2] on convex sets are also substantially extended to star-shaped sets.

Let  $X$  be a complete metric space and  $K \subset X$ . We say that a map  $T : K \rightarrow X$  is metrically inward if for each  $x \in K$  there exists an element  $u$  of  $K$  such that  $d(x, u) + d(u, Tx) = d(x, Tx)$  where  $u = x$  iff  $x = Tx$  [7].

Let  $x_0 \in K$ . We shall construct two sequences  $\{x_n\}$ ,  $\{x_n'\}$  as follows: Define  $x_1' = Tx_0$ . If  $x_1' \in K$ , set  $x_1 = x_1'$ . If  $x_1' \notin K$ , choose  $x_1 \in K$  so that  $d(x_0, x_1) + d(x_1, x_1') = d(x_0, x_1')$ . Set  $x_2' = Tx_1$ . If  $x_2' \in K$ , set  $x_2 = x_2'$ . If not, choose  $x_2 \in K$  so that  $d(x_1, x_2) + d(x_2, x_2') = d(x_1, x_2')$ . Continuing in this manner, we obtain  $\{x_n\}$ ,  $\{x_n'\}$  satisfying

- (1)  $x_{n+1}' = Tx_n$ ,
- (2)  $x_n = x_n'$  if  $x_n' \in K$ , and
- (3)  $x_n \in K$  and  $d(x_{n-1}, x_n) + d(x_n, x_n') = d(x_{n-1}, x_n')$  if  $x_n' \notin K$ .

Let  $P = \{x_i \in \{x_n\} \mid x_i = x_i'\}$  and  $Q = \{x_i \in \{x_n\} \mid x_i \neq x_i'\}$ .

The following is our main result which is comparable to Theorem 2.2 of Caristi [7].

**THEOREM 1.** *Let  $(X, d)$  be a complete metric space,  $K$  a nonempty closed subset of  $X$ , and  $T : K \rightarrow X$  a metrically inward map satisfying the condition:*

- (A) *given  $\varepsilon > 0$ , there exists a  $\delta > 0$  such that  $x, y \in K$  and*

$$\varepsilon \leq d(x, y) < \varepsilon + \delta \text{ imply } d(Tx, Ty) < \varepsilon.$$

*If a sequence  $\{x_n\}$  defined as above satisfies the condition:*

---

Received Sept. 24, 1981.

\*Supported by a grant of the Korea Science and Engineering Foundation, 1980-1981.

(\*)  $x_n \in Q$  implies  $x_{n-1}, x_{n+1} \in P$ ,

then  $T$  has a unique fixed point and  $\{x_n\}$  converges to the fixed point.

*Proof.* If there exists a  $j$  such that  $x_n \in P$  for all  $n \geq j$ , then  $\{x_n\}$  converges to a fixed point of  $T$  by the method of Meir-Keeler [13]. Therefore, we may assume that  $Q$  contains infinitely many elements. Let  $Q = \{x_{n_k}\}$ . If  $x_j = x_{j+1}$  for some  $j$ , then  $x_j = Tx_j$  is a fixed point. Therefore, we may assume  $x_n \neq x_{n+1}$  and hence  $x_n \neq x_{n+1}'$  for all  $n$ .

Step 1.  $d(x_{n_k-1}, x_{n_k}') \rightarrow 0$  as  $k \rightarrow \infty$ .

Set  $n_k = r$  and  $n_{k+1} = s$ . Since  $T$  is contractive and  $r < s-1$ , we have

$$\begin{aligned} d(x_{s-1}, x_s') &= d(Tx_{s-2}, Tx_{s-1}) < d(x_{s-2}, x_{s-1}) < \dots \\ &< d(x_r, x_{r+1}) \leq d(x_r, x_r') + d(x_r', x_{r+1}) \\ &= d(x_r, x_r') + d(Tx_{r-1}, Tx_r) \\ &< d(x_r, x_r') + d(x_{r-1}, x_r) \\ &= d(x_{r-1}, x_r'). \end{aligned}$$

Therefore  $\{d(x_{r-1}, x_r')\}$  is decreasing to a limit  $\varepsilon \geq 0$ . Suppose  $\varepsilon > 0$ . Then there exists a  $\delta > 0$  satisfying (A), and hence there exists an  $N$  such that  $\varepsilon \leq d(x_{n_k-1}, x_{n_k}') < \varepsilon + \delta$  for all  $k \geq N$ . Since  $d(x_{s-1}, x_s') < d(x_r, x_{r+1}) < d(x_{r-1}, x_r')$ , we have  $\varepsilon \leq d(x_r, x_{r+1}) < \varepsilon + \delta$ , and hence  $d(Tx_r, Tx_{r+1}) < \varepsilon$  by (A). On the other hand,

$$\begin{aligned} d(x_{s-1}, x_s') &= d(Tx_{s-2}, Tx_{s-1}) < d(x_{s-2}, x_{s-1}) = d(x_{s-2}, x_{s-1}') \\ &< \dots < d(x_{r+1}, x_{r+2}') = d(Tx_r, Tx_{r+1}) < \varepsilon, \end{aligned}$$

which is a contradiction.

Step 2.  $d(x_n, x_{n+1}) \rightarrow 0$ ,  $d(x_n, Tx_n) \rightarrow 0$  as  $n \rightarrow \infty$ .

Suppose  $d(x_n, x_{n+1}) \rightarrow 0$  is false. Then there exists an  $\varepsilon > 0$  such that for every  $N \geq 0$ , there exists an  $n \geq N$  such that  $d(x_{n-1}, x_n) > \varepsilon$ . By Step 1, there exists an  $M \geq 0$  such that  $d(x_{n_k-1}, x_{n_k}') < \varepsilon$  for  $k \geq M$ . Let  $N = n_k$  for some  $k \geq M$ . Then for all  $n \geq N$  such that  $x_n \in P$ , there exists a unique  $j$  such that  $n_j < n < n_{j+1}$ , and hence

$$\begin{aligned} d(x_{n-1}, x_n) &= d(x_{n-1}, x_n') = d(Tx_{n-2}, Tx_{n-1}) < d(x_{n-2}, x_{n-1}) \\ &= d(x_{n-2}, x_{n-1}') < \dots < d(x_{n_j-1}, x_{n_j}') < \varepsilon, \end{aligned}$$

a contradiction. Similar argument shows that  $d(x_n, Tx_n) \rightarrow 0$ .

Step 3.  $\{x_n\}$  is Cauchy.

Suppose not. Then there exists an  $\varepsilon > 0$  such that  $\limsup_{m,n} d(x_m, x_n) > 2\varepsilon$ . Now there exists a  $\delta$ ,  $0 < \delta \leq \varepsilon$ , satisfying (A). By Step 2, there exists an  $M$  such that for  $n \geq M$ ,

$$d(x_n, x_{n+1}) < \delta/3 \text{ and } d(x_n, Tx_n) < \delta/3.$$

Choose  $m > n > M$  so that  $d(x_m, x_n) > 2\varepsilon$ . For  $j$ ,  $m \geq j \geq n$ ,

$$d(x_m, x_j) \leq d(x_m, x_{j+1}) + d(x_j, x_{j+1})$$

implies

$$|d(x_m, x_j) - d(x_m, x_{j+1})| \leq d(x_j, x_{j+1}) < \delta/3.$$

Since  $d(x_m, x_{m+1}) < \delta/3 < \varepsilon$  and  $d(x_m, x_n) > 2\varepsilon \geq \varepsilon + \delta$ , there exists a  $j$ ,  $m \geq j \geq n$ , such that  $\varepsilon + 2\delta/3 < d(x_m, x_j) < \varepsilon + \delta$ . On the other hand,

$$\begin{aligned} d(x_m, x_j) &\leq d(x_m, Tx_m) + d(Tx_m, Tx_j) + d(Tx_j, x_j) \\ &< \delta/3 + \varepsilon + \delta/3 = \varepsilon + 2\delta/3, \end{aligned}$$

a contradiction.

Since  $\{x_n\}$  is Cauchy, it converges to some  $p \in K$ . Since  $d(x_n, Tx_n) \rightarrow 0$ , we have  $Tx_n \rightarrow p$  and hence  $p = Tp$  by the continuity of  $T$ . The uniqueness is clear. This completes our proof.

REMARK 1.1. The condition (A) is due to Meir-Keeler [13]. Since every selfmap is metrically inward and satisfies (\*), Theorem 1 extends Meir-Keeler's theorem [13]. Note that the condition (A) includes the following:

- (A<sub>1</sub>) given  $\varepsilon > 0$ , there exists a  $\delta > 0$  and an  $\varepsilon_0$  with  $0 < \varepsilon_0 < \varepsilon$  such that  $\varepsilon \leq d(x, y) < \varepsilon + \delta$  implies  $d(Tx, Ty) \leq \varepsilon_0$ ;
- (A<sub>2</sub>) given  $\varepsilon > 0$ , there exists a  $\delta > 0$  and an  $\varepsilon_0$  with  $0 < \varepsilon_0 < \varepsilon$  such that  $d(x, y) < \varepsilon + \delta$  implies  $d(Tx, Ty) \leq \varepsilon_0$ ;
- (A<sub>3</sub>)  $T$  is a Banach contraction, that is, there exists a  $k < 1$  such that  $d(Tx, Ty) \leq k d(x, y)$ .

Note that (A<sub>3</sub>)  $\Rightarrow$  (A<sub>2</sub>)  $\Rightarrow$  (A<sub>1</sub>)  $\Rightarrow$  (A). Hegedüs-Szilágyi [9] noted that (A<sub>1</sub>) is equivalent to the contractive type condition of Boyd-Wong [5], and (A<sub>2</sub>) to the condition of Browder [6].

REMARK 1.2. Caristi ([4], Theorem 2.2) obtained Theorem 1 for the condition (A<sub>3</sub>) instead of (A) without assuming the condition (\*). He showed merely the existence of fixed point using the Caristi-Kirk fixed point theorem.

The star-shapedness of linear spaces [11] can be extended for metric spaces as follows:

A metric space  $(X, d)$  is said to be star-shaped if there exists at least one point  $c \in X$  (called a star-center) such that for each  $x \in X$ ,  $c$  and  $x$  can be

joined by a metric segment of  $X$ , that is, a subset isometric to an interval of length  $d(c, x)$ . A metric space  $X$  is said to be star-shaped with respect to a subset  $K$  of  $X$  if each  $y \in K$  is a star-center of  $X$

REMARK 1.3. A metric space  $X$  is convex in the sense of Bing [4] if for every  $x, y \in X$  there exists a  $z \in X$  such that  $d(x, y)/2 = d(x, z) = d(z, y)$ , and convex in the sense of Menger if for every  $x, y \in X$  there exists a  $z \in X$  such that  $d(x, y) = d(x, z) + d(z, y)$ . Clearly, the former implies the latter. However, for complete metric spaces those concepts are identical since a theorem of Menger (see [10]) states that a convex complete metric space contains together with  $x, y$  also a metric segment whose extremities are  $x$  and  $y$ .

REMARK 1.4. Matkowski and Wegrzyk [12] claimed that on a complete, metrically convex, metric space, the conditions (A),  $(A_1)$ ,  $(A_2)$  and some others are equivalent.

In certain cases, while constructing the sequences  $\{x_n\}, \{x_n'\}$  as in Theorem 1, if  $x_n' \notin K$ ,  $x_n$  can be chosen at the boundary of  $K$ . From now on,  $\partial$  denotes the boundary and  $\partial_X$  the relative boundary.

THEOREM 2. *Let  $(X, d)$  be a complete metric space and  $K$  a closed subset of  $X$  such that  $X$  is star-shaped with respect to  $K$ . If a map  $T: K \rightarrow X$  satisfies the condition (A) and  $T(\partial K) \subset K$ , then  $T$  has a unique fixed point and a sequence  $\{x_n\}$  defined as above converges to the fixed point.*

*Proof.* Since  $X$  is star-shaped with respect to  $K$ ,  $T$  is metrically inward. Note also that the assumption  $T(\partial K) \subset K$  implies (\*). Therefore Theorem 2 follows from Theorem 1.

COROLLARY 2.1. (Assad [3]) *Let  $(X, d)$  be a complete, metrically convex, metric space and  $K$  a nonempty closed subset of  $X$ . Suppose that  $T: K \rightarrow X$  satisfies (A) and  $T(\partial K) \subset K$ . Then  $T$  has a unique fixed point.*

COROLLARY 2.2. *Let  $E$  be a Banach space,  $X$  a closed subset of  $E$  and,  $K$  a closed subset of  $X$  such that  $X$  is star-shaped with respect to  $K$ . If  $T: K \rightarrow X$  satisfies the condition (A) and  $T(\partial_X K) \subset K$ , then  $T$  has a unique fixed point.*

REMARK 2.1. Kuhfittig [11] obtained Corollary 2.2 for the condition  $(A_2)$  instead of (A), and Assad-Kirk [1] for the condition  $(A_3)$  and convex  $X$ .

Following Dotson's method in [8], from Corollary 2.2 we obtain

COROLLARY 2.3. (Kuhfittig [11]) *Let  $E$  be a Banach space,  $X$  a closed subset of  $E$ , and  $K$  a compact star-shaped subset of  $X$  such that  $X$  is star-shaped with respect to  $K$ . If  $T: K \rightarrow X$  is nonexpansive, and if  $T(\partial_X K) \subset K$ , then  $T$  has a fixed point.*

REMARK 2.2. Corollary 2.3 extends Theorem 1 of Dotson [8]. Note that other main results in [11] follows from Corollary 2.2.

The following is closely related to Theorem 1 and its proof can be given by modifying that of Rhoades' theorem [14].

THEOREM 3. Let  $(X, d)$  be a complete metric space,  $K$  a closed subset of  $X$ , and  $T : K \rightarrow X$  a metrically inward map satisfying the condition:

(B) there exists an  $h \in (0, 1)$  such that for each  $x, y \in K$ ,  
 $d(Tx, Ty) \leq h \max \{d(x, y)/2, d(x, Tx), d(y, Ty), [d(x, Ty) + d(y, Tx)]/q\}$   
 where  $q \in [1 + 2h, \infty)$ . If a sequence  $\{x_n\}$  defined as in Theorem 1 satisfies the condition (\*), then  $T$  has a unique fixed point and  $\{x_n\}$  converges to the point.

COROLLARY 3.1. Let  $(X, d)$  be a complete metric space and  $K$  a closed subset of  $X$  such that  $X$  is star-shaped with respect to  $K$ . If a map  $T : K \rightarrow X$  satisfies the condition (B) and  $T(\partial K) \subset K$ , then  $T$  has a unique fixed point.

COROLLARY 3.2. Let  $E$  be a Banach space,  $X$  a closed subset of  $E$ , and  $K$  a closed subset of  $X$  such that  $X$  is star-shaped with respect to  $K$ . If  $T : K \rightarrow X$  satisfies the condition (B) and  $T(\partial_X K) \subset K$ , then  $T$  has a unique fixed point in  $K$ .

REMARK 3. Rhoades [14] obtained Corollary 3.2 for convex  $X$ , and Assad [2] for  $T$  satisfying the condition

$$(B_1) \quad \|Tx - Ty\| \leq k(\|x - Tx\| + \|y - Ty\|), \quad k < 1/2.$$

Note that  $(B_1)$  implies (B).

## References

1. N. A. Assad and W. A. Kirk, *Fixed point theorems for set-valued mappings of contractive type*, Pacific J. Math. **43** (1972), 553-562.
2. ———, *On a fixed point theorem of Kannan in Banach spaces*, Tamkang J. Math. **7** (1976), 91-94.
3. ———, *A fixed point theorem for weakly uniformly strict contractions*, Canad. Math. Bull. **16** (1973), 15-18.
4. R. H. Bing, *A convex metric for a locally connected continuum*, Bull. Amer. Math. Soc. **55** (1949), 812-819.
5. D. W. Boyd and J. S. W. Wong, *On nonlinear contractions*, Proc. Amer. Math. Soc. **20** (1969), 458-464.
6. F. E. Browder, *On the convergence of successive approximations for nonlinear functional equations*, Indag. Math. **30** (1968), 27-35.

7. J. Caristi, *Fixed point theorems for mappings satisfying inwardness conditions*, Trans. Amer. Math. Soc. **215** (1976), 241-251.
8. W.G. Dotson, *Fixed point theorems for non-expansive mappings on star-shaped subsets of Banach spaces*, J. London Math. Soc. (2) **4** (1972), 408-410.
9. M. Hegedüs and T. Szilágyi, *Equivalent conditions and a new fixed point theorem in the theory of contractive type mappings*, Math. Japonica **25** (1980), 147-157.
10. W. A. Kirk, *Caristi's fixed point theorem and metric convexity*, Colloq. Math. **36** (1976), 81-86.
11. P. Kuhfittig, *Fixed point theorems for mappings with non-convex domain and range*, Rocky Mountain J. Math. **7** (1977), 141-145.
12. J. Matkowski and R. Wegrzyk, *On equivalence of some fixed point theorems for selfmappings of metrically convex spaces*, Bollettino U.M.I. (5) **15** (1978), 359-369.
13. A. Meir and E. Keeler, *A theorem on contraction mappings*, Jour. Math. Anal. Appl. **28** (1969), 326-329.
14. B. E. Rhoades, *A fixed point theorem for some non-self-mappings*, Math. Japonica **23** (1978), 457-459.

Seoul National University