

Research Article

Remarks on Extensions of the Himmelberg Fixed Point Theorem

Hidetoshi Komiya and Sehie Park

Received 30 August 2007; Accepted 16 November 2007

Recommended by Anthony To-Ming Lau

Recently, Jafari and Sehgal obtained an extension of the Himmelberg fixed point theorem based on the Kakutani fixed-point theorem. We give generalizations of the extension to almost convex sets instead of convex sets. We also give generalizations for a large class \mathfrak{B} of better admissible multimaps instead of the Kakutani maps. Our arguments are based on the KKM principle and some of previous results due to the second author.

Copyright © 2007 H. Komiya and S. Park. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In 1972, Himmelberg [1] derived the following from the Kakutani fixed point theorem.

THEOREM 1.1. *Let T be a nonvoid convex subset of a separated locally convex space L . Let $F : T \rightarrow T$ be a u.s.c. multimap such that $F(x)$ is closed and convex for all $x \in T$, and $F(T)$ is contained in some compact subset C of T . Then F has a fixed point.*

Recall that Theorem 1.1 is usually called *the Himmelberg fixed point theorem* and is a common generalization of historically well-known fixed point theorems due to Brouwer, Schauder, Tychonoff, Kakutani, Bohnenblust and Karlin, Fan, Glicksberg, and Hukuhara (see [2]). Recall also that the multimap F is usually called a *Kakutani map*.

Recently, Jafari and Sehgal [3] obtained an extension of Theorem 1.1 based on the Kakutani fixed point theorem. Our aim in this paper is to give generalizations of the extension to almost convex sets instead of convex sets. We also give generalizations for a large class \mathfrak{B} of better admissible multimaps instead of the Kakutani maps. Our arguments are based on the KKM principle and some results in [3–5].

2 Fixed Point Theory and Applications

2. Preliminaries

Recall that, for topological spaces X and Y , a multimap (simply, a map) $F : X \multimap Y$ is u.s.c. (resp., l.s.c.) if, for any closed (resp., open) subset $A \subset X$,

$$F^{-1}(A) := \{x \in X \mid F(x) \cap A \neq \emptyset\} \quad (2.1)$$

is closed (resp., open) in X . If Y is regular, F is u.s.c. and has nonempty closed values, then F has a closed graph.

Himmelberg [1] defined that a subset A of a t.v.s. E is said to be *almost convex* if, for any neighborhood V of the origin 0 in E and for any finite set $\{w_1, \dots, w_n\}$ of points of A , there exist $z_1, \dots, z_n \in A$ such that $z_i - w_i \in V$ for all i , and $\text{co}\{z_1, \dots, z_n\} \subset A$.

As the second author once showed in [6], the classical KKM principle implies many fixed point theorems. In [4], the following almost fixed point theorem was obtained from the KKM principle.

THEOREM 2.1. *Let X be a subset of a t.v.s. and Y an almost convex dense subset of X . Let $T : X \multimap E$ be an l.s.c. (resp., a u.s.c.) map such that $T(y)$ is convex for all $y \in Y$. If there is a totally bounded subset K of \bar{X} such that $T(y) \cap K \neq \emptyset$ for each $y \in Y$, then for any convex neighborhood V of the origin 0 of E , there exists a point $x_V \in Y$ such that $T(x_V) \cap (x_V + V) \neq \emptyset$.*

Note that a t.v.s. is not necessarily Hausdorff in Theorem 2.1. It is routine to deduce Theorem 1.1 from Theorem 2.1. In fact, in 2000, we had the following in [7].

THEOREM 2.2. *Let X be a subset of a locally convex Hausdorff t.v.s. E and Y an almost convex dense subset of X . Let $T : X \multimap X$ be a compact u.s.c. map with nonempty closed values such that $T(y)$ is convex for all $y \in Y$. Then T has a fixed point.*

In particular, for $Y = X$, we obtain the following generalization [7] of Theorem 1.1.

THEOREM 2.3. *Let X be an almost convex subset of a locally convex Hausdorff t.v.s. Then any compact u.s.c. map $T : X \multimap X$ with nonempty closed convex values has a fixed point in X .*

A *polytope* P in a subset X of a t.v.s. E is a subset of X homeomorphic to a standard simplex.

We define “better” admissible class \mathfrak{B} of maps from a subset X of a t.v.s. E into a topological space Y as follows.

$F \in \mathfrak{B}(X, Y) \Leftrightarrow F : X \multimap Y$ is a map such that, for each polytope P in X and for any continuous function $f : F(P) \rightarrow P$, the composition $f(F|_P) : P \rightarrow P$ has a fixed point.

There is a large number of examples of better admissible maps (see [5]). A typical example is an *acyclic map*, that is, a u.s.c. map with compact acyclic values. It is also known that any u.s.c. map with compact values having a *trivial shape* (i.e., contractible in each neighborhood) belongs to $\mathfrak{B}(X, Y)$, see [8].

For a subset C of a t.v.s. E , we say that a multimap $F : C \multimap C$ has an *E -almost fixed point* if, for each neighborhood V of the origin 0 in E , there exist points $x_V \in C$ and $y_V \in F(x_V)$ such that $x_V - y_V \in V$ as in Theorem 2.1.

The following generalization of Theorems 1.1 and 2.3 is a consequence of the main theorem of [5], where \mathfrak{B}^p should be replaced by \mathfrak{B} .

THEOREM 2.4. *Let X be an almost convex subset of a locally convex t.v.s. E .*

- (1) *If $F \in \mathfrak{B}(X, X)$ is compact, then F has an E -almost fixed point.*
- (2) *Further, if E is Hausdorff and F is closed, then F has a fixed point.*

In what follows, let $E = (E, \tau)$ be a t.v.s. with topology τ , $\hat{E} = (E, \tau)^\wedge$ the completion of E , and E^* the topological dual of E . Recall that if E^* separates points of E , then (E, τ) is Hausdorff and (E, τ_w) with the weak topology is Hausdorff and locally convex. We will use the following lemmas in [3].

LEMMA 2.5 [3, Lemma 2]. *If $(\hat{E})^*$ separates points of \hat{E} , then E^* separates points of E .*

LEMMA 2.6 [3, Lemma 4]. *Let $(\hat{E})^*$ separate points of \hat{E} . Let K be a compact subset of E whose $\overline{\text{co}}K$ in \hat{E} is \hat{E} -compact. If a net $\{x_\alpha\} \subset \text{co}K$ is such that for some $u \in K$, $\{x_\alpha\} \rightarrow u$ in (E, τ_w) , then there is a subnet $\{x_\beta\}$ of $\{x_\alpha\}$ with $\{x_\beta\} \rightarrow u$ in (E, τ) .*

3. New fixed point theorems

Motivated by [3], we obtain the following main result of this paper.

THEOREM 3.1. *Let E be a t.v.s., C an almost convex subset of E , and K a compact subset of C such that $\overline{\text{co}}K$ is \hat{E} -compact. Let $F : C \multimap K$ be a u.s.c. multimap such that*

- (1) *for each $x \in C$, $F(x)$ is a nonempty closed subset of K ;*
- (2) *F has an (E, τ_w) -almost fixed point in K .*

If $(\hat{E}, \tau_w)^$ separates points of \hat{E} , then F has a fixed point in K .*

Proof. We follow that of [3, Theorem 5]. Since $(\hat{E}, \tau_w)^*$ separates points of \hat{E} , by Lemma 2.5, (E, τ_w) is a Hausdorff locally convex t.v.s. Let \mathcal{U} be a neighborhood basis of the origin 0 of (E, τ_w) consisting of (E, τ_w) -closed convex and symmetric subsets of E . For each $V \in \mathcal{U}$, there exist points $x_V \in K$, $y_V \in F(x_V)$ such that $x_V - y_V \in V$. Partially order \mathcal{U} by inclusion. Then $\{x_V - y_V \mid V \in \mathcal{U}\} \rightarrow 0$ in (E, τ_w) . Since $\{y_V \mid V \in \mathcal{U}\} \subset K$, there exists a subnet $\{y_{V'} \mid V' \in \mathcal{U}' \subset \mathcal{U}\}$ and a $u \in K$ such that $\{y_{V'} \mid V' \in \mathcal{U}'\} \rightarrow u$ in E . Since $x_V - y_V \in V$, the net $\{x_{V'} \mid V' \in \mathcal{U}'\} \rightarrow u$ in (E, τ_w) . Since $u \in K$ and $\{x_{V'} \mid V' \in \mathcal{U}'\} \subset \text{co}K$, it follows by Lemma 2.6 that there is a subnet $\{x_{V''}\}$ of $\{x_{V'}\}$ with $\{x_{V''}\} \rightarrow u$ in E . Hence $\{y_{V''}\} \rightarrow u$ in E also. Since K is regular and F is u.s.c. with closed values, F has a closed graph. Since, for each V'' , $y_{V''} \in F(x_{V''})$, we have $u \in F(u)$. This completes our proof. □

From Theorem 2.1, we immediately have the following.

THEOREM 3.2. *Let E be a t.v.s. such that (E, τ_w) is a locally convex t.v.s., C an almost convex subset of E , and K a (E, τ_w) -totally bounded subset of C . Let $F : C \multimap K$ be a u.s.c. (resp., an l.s.c.) multimap such that for each $x \in C$, $F(x)$ is a nonempty convex subset of K . Then F has an (E, τ_w) -almost fixed point in K .*

4 Fixed Point Theory and Applications

Combining Theorems 3.1 and 3.2, we have the following.

THEOREM 3.3. *Let E be a t.v.s., C an almost convex subset of E , and K a compact subset of C such that $\overline{co}K$ is \hat{E} -compact. Let $F : C \multimap K$ be a u.s.c. multimap such that for each $x \in C$, $F(x)$ is a nonempty closed and convex subset of K . If $(\hat{E}, \tau_w)^*$ separates points of \hat{E} , then F has a fixed point in K .*

When C is convex, Theorem 3.3 reduces to the main theorem of Jafari and Sehgal [3]. As noted in [3], if E is a locally convex Hausdorff t.v.s., then so is \hat{E} and hence $(\hat{E})^*$ separates points of \hat{E} . Consequently, Theorem 2.2 follows from Theorem 3.3.

From Theorem 2.4, we immediately have the following.

THEOREM 3.4. *Let E be a t.v.s. such that (E, τ_w) is a locally convex t.v.s., C an almost convex subset of E , and K an (E, τ_w) -compact subset of C . If $F \in \mathfrak{B}(C, K)$, then F has an (E, τ_w) -almost fixed point in K .*

Combining Theorems 3.1 and 3.4, we have the following.

THEOREM 3.5. *Let E be a t.v.s., C an almost convex subset of E , and K a compact subset of C such that $\overline{co}K$ is \hat{E} -compact. Let $F \in \mathfrak{B}(C, K)$. If $(\hat{E}, \tau_w)^*$ separates points of \hat{E} , then F has a fixed point in K .*

As noted in [3], if E is a locally convex Hausdorff t.v.s., then so is \hat{E} and hence $(\hat{E})^*$ separates points of \hat{E} . Consequently, Theorem 2.4 follows from Theorem 3.5.

Acknowledgments

This work was done while the second author was visiting Keio University in the Fall, 2006. He would like to express his gratitude to the university authorities for their kind hospitality.

References

- [1] C. J. Himmelberg, "Fixed points of compact multifunctions," *Journal of Mathematical Analysis and Applications*, vol. 38, pp. 205–207, 1972.
- [2] S. Park, "Ninety years of the Brouwer fixed point theorem," *Vietnam Journal of Mathematics*, vol. 27, no. 3, pp. 187–222, 1999.
- [3] F. Jafari and V. M. Sehgal, "An extension to a theorem of Himmelberg," *Journal of Mathematical Analysis and Applications*, vol. 327, no. 1, pp. 298–301, 2007.
- [4] S. Park, "The Knaster-Kuratowski-Mazurkiewicz theorem and almost fixed points," *Topological Methods in Nonlinear Analysis*, vol. 16, no. 1, pp. 195–200, 2000.
- [5] S. Park, "Fixed point theorems for better admissible multimaps on almost convex sets," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 1, pp. 690–702, 2007.
- [6] S. Park, "The KKM principle implies many fixed point theorems," *Topology and Its Applications*, vol. 135, no. 1–3, pp. 197–206, 2004.
- [7] S. Park and D. H. Tan, "Remarks on Himmelberg-Idzik's fixed point theorem," *Acta Mathematica Vietnamica*, vol. 25, no. 3, pp. 285–289, 2000.

- [8] H. Ben-El-Mechaiekh, “Spaces and maps approximation and fixed points,” *Journal of Computational and Applied Mathematics*, vol. 113, no. 1-2, pp. 283–308, 2000.

Hidetoshi Komiya: Faculty of Business and Commerce, Keio University, Hiyoshi,
Yokohama 223-8521, Japan


Email address: hkomiya@fbc.keio.ac.jp

Sehie Park: The National Academy of Sciences, Seocho-gu, Seoul 137-044, Korea; Department of
Mathematical Sciences, College of Natural Science, Seoul National University, Seoul 151-747, Korea

Email address: shpark@math.snu.ac.kr

Impulsive Differential Equations and Inclusions

M. Benchohra, J. Henderson, and S. Ntouyas



Functional differential equations and inclusions have assumed an increasing relevance, mainly due to the great variety of problems arising in sciences, engineering, and technology which can be described by such types of equations and inclusions. This book is devoted to this rapidly developing theory of differential equations and inclusions with special emphasis on such equations and inclusions subject to impulses. In addition, the book includes a wide account of various applications to physical problems.

Limited-Time Promotional Offer. Buy this title NOW at **20% discount plus Free Shipping.**

The book consists of twelve chapters. Initial value problems and boundary value problems for both impulsive differential equations and differential inclusions, as well as for both impulsive functional differential equations and functional differential inclusions, or for both neutral functional differential equations and neutral functional differential

inclusions, are studied. Moreover, results on impulsive semilinear ordinary differential inclusions and functional differential inclusions satisfying nonlocal boundary conditions are given extensive attention. Positive solutions and multiple positive solutions for impulsive ordinary differential equations and functional differential equations, as well as impulsive differential inclusions satisfying periodic boundary conditions, are also studied.

Some of these results are extended to functional differential equations and functional differential inclusions with variable times, to semilinear functional differential equations and functional differential inclusions for operators that are nondensely defined on a Banach space, to impulsive hyperbolic differential inclusions, and to impulsive dynamic equations on time scales.

Each chapter concludes with a section devoted to notes and bibliographical remarks. This book is addressed to a wide audience of specialists such as mathematicians, engineers, biologists, and physicists.

CMIA Book Series, Volume 2, ISBN: 977-5945-50-X

Please visit <http://www.hindawi.com/cmia.2.html> for more information about the book.

To place an order while taking advantage of our current promotional offer, please contact books.orders@hindawi.com